

**Evaluating Ecosystem Management Capabilities at the Local Level in Florida:
A Policy Gap Analysis Using Geographic Information Systems**

Samuel D. Brody*
Assistant Professor
Department of Landscape Architecture and Urban Planning
Texas A&M University
TAMU 3137
College Station, Texas 77843-3137
Phone: (979) 458-4623 Email: sbrody@archone.tamu.edu

Virginia Carrasco
Center for Urban Housing and Development Graduate Fellow
Department of Landscape Architecture and Urban Planning
Texas A&M University
TAMU 3137
College Station, Texas 77843-3137

Wes Highfield
Sustainable Coastal Margins Program Graduate Fellow
Department of Landscape Architecture and Urban Planning
Texas A&M University
TAMU 3137
College Station, Texas 77843-3137

February 2002

** Corresponding author*

Evaluating Ecosystem Management Capabilities

Abstract

Because ecosystem approaches to management adhere to ecological systems rather than human-defined boundaries, collaboration across jurisdictions, agencies, and land ownership is often necessary to achieve effective management of transboundary resources. Local natural resource and land use planners increasingly recognize that while ecosystem management requires looking beyond specific jurisdictions and focusing on broad spatial scales, the approach will partly be implemented at the local level with the coordination of local policies across larger landscapes. This article evaluates the collective capabilities of local jurisdictions to manage large transboundary ecological systems in Florida. Specifically, it combines plan evaluation with Geographic Information Systems (GIS) techniques to map, measure, and analyze the existing mosaic of management across selected ecosystems in the southern portion of the State. Visual and statistical results indicate significant gaps in the management framework of southern Florida that, if filled, could achieve a greater level of consistency and more complete coverage of ecosystem management policies. Based on the findings, we make recommendations on how and where local plans can be strengthened to more effectively attain the objectives of ecosystem approaches to management.

Key Words: ecosystem management; collaboration; Geographic Information Systems; land use policies

Evaluating Ecosystem Management Capabilities

In the United States (U.S.), environmental planners are abandoning the traditional species-by-species approach to regulation where policies are fragmented by administrative lines, and instead are embracing the emerging practice of ecosystem management. Ecosystem management represents a departure from traditional types of management by addressing the interaction between biotic and abiotic components within a landscape, while incorporating human concerns (Szaro *et al.* 1998). In this approach, entire ecological systems (e.g. watersheds, ecological communities, etc.), and the ecological structures, functions, and processes within them, become the focus for management efforts rather than a single species or jurisdiction (Grumbine 1994; Vogt *et al.* 1997). An ecosystem management framework is unique because it recognizes the connections to large-scale natural processes that extend beyond individual ownership, management jurisdictions, and political boundaries (Christensen *et al.* 1996).

Ecosystem approaches to management thus attempt to transcend jurisdiction lines by broadening managers' geographic focus and creating situations of collaborative problem solving. This management framework is based on an ecosystem science that integrates many disciplinary approaches and addresses ecological issues often at very large temporal and spatial scales (Peck 1998). By aligning policies and plans with a coherent, ecologically defined spatial unit, ecosystem approaches to management can more effectively protect ecological structure, function, and overall biodiversity.

Evaluating Ecosystem Management Capabilities

Local natural resource and land use planners increasingly recognize that while ecosystem management requires looking beyond specific jurisdictions and focusing on broad spatial scales, the approach will in part be implemented at the local level through local land use decisions. Furthermore, ecosystem approaches to management may not be realized solely by structural or engineering approaches to management, but by the coordination of local plans and policies across larger landscapes (Kirklin 1995; Beatley 2000). Local level planning therefore must be considered along with other spatial and jurisdictional scales when it comes to managing entire ecological systems. The factors causing ecosystem decline, such as rapid urban development and habitat fragmentation occur at the local level and are generated by local land use decisions (Noss and Scott 1997). The vast majority of these decisions affecting large ecosystems will be made at a smaller scale where they make the largest impact on the natural environment (Endter-Wada 1998; McGinnis *et al.* 1999). As a result, many of the decisions which may threaten or protect natural habitat are in the hands of county commissioners, city councils, town boards, local planning staff, and the participating public. Thoughtful policies and actions at the local level can often protect critical habitats of regional significance more effectively and less expensively than the best intentioned state or federal protection schemes (Duerksen *et al.* 1997). The importance of local ecosystem-based planning is further highlighted by the declining role of the federal government in the protection of habitat and associated ecological systems over the past ten years, and a future political climate that suggests giving more control to local jurisdictions for making resource use decisions.

While a large amount of research has been conducted on instituting the broad principles of ecosystem management, comparatively little work has been done to show how natural systems

Evaluating Ecosystem Management Capabilities

can be effectively managed at the local level. This article evaluates the collective capabilities of local jurisdictions to manage large transboundary ecological systems. Specifically, it uses Geographic Information Systems (GIS) to map, measure, and analyze the existing mosaic of management based on policies in comprehensive plans across selected ecosystems in the southern portion of Florida. Our study seeks to provide answers to the following questions: 1) what is the degree of spatial coverage of ecosystem strategies and policies in southern Florida; 2) what is the existing spatial pattern of select indicators within specific ecological systems; 3) how well are multiple local jurisdiction collectively managing larger ecosystems; and 4) are plan scores randomly distributed across the study area, or clustered within particular ecological units?

The following section examines the importance of collaboration among organizations and jurisdictions in managing ecological systems, specifically in south Florida. Based on the defining principles of collaborative ecosystem management, we derive a set of indicators to assess the ability of local comprehensive plans to manage natural systems extending beyond a single jurisdiction. Next, sample selection, variable measurement, and data analysis procedures are described. Visual and statistical results indicate the degree of coverage for single and aggregated policies across specific ecosystems and pinpoint gaps that can be filled by future plan updates to obtain a more consistent management framework across the State. Descriptive results identify policy gaps in the framework of management and increase understanding of how and exactly where to incorporate ecosystem management most effectively into local level resource planning decisions. By using GIS-based methods, the findings provide guidance for planners on where to establish future land use policies to reduce the decline of biological diversity and protect critical ecosystem components. Finally, based on the ecosystem management gap

analysis, policy implications are discussed and suggestions are made as to how and where land use management can be strengthened by future planning initiatives.

The Importance of Collaboration In Ecosystem Approaches To Management

Ecosystem approaches to management increasingly depend on collaboration across political, administrative, and ownership boundaries (Blumenthal and Jannink 2000; Selin *et al.* 2000). Because these approaches adhere to ecological systems rather than administrative or political lines, inter-organizational collaboration across jurisdictions, agencies, and land ownership is often necessary to achieve effective management of transboundary resources. Management decisions must be made collectively because in most cases no single entity has jurisdiction over all aspects of an ecosystem. The need to integrate the values and knowledge of a broad array of organizations and individuals translates into a need to focus on collaborative planning efforts among resource owners, managers, and users (Cortner and Moote 1999; Wondolleck and Yaffee 2000).

In recent years, the organizational design literature has increasingly called for inter-organizational and intersectoral collaboration to solve major environmental problems (Gray 1989; Cooperider and Passmore 1991). Such collaborations, which include a variety of inter-jurisdictional partnerships and public-private alliances, have been viewed as critical to effective management outcomes which meet the needs of all interested parties (Westley and Vrendenburg 1997). In this sense, collaborations induced by shared visions are intended to advance the collective good of the stakeholders involved (Bryson and Crosby 1992). Emery and Trist (1965)

initially argued that problem domains (ill- defined problems that depend on multiple perspectives for their solution) could be stabilized by inter-organizational collaboration. These so called “meta-problems” which transcend boundaries of single organizations or jurisdictions must be addressed cooperatively. Others have also noted that multiple stakeholders in different sectors, having different viewpoints, interests, and values, must cooperate to solve problems whose parameters are transboundary or not clearly defined (Clark 1989; Brown 1991). In an extensive review of the nature of collaborative alliances, Gray and Wood (1991) found that many organizations, particularly those from the private sector, are finding it advantageous and often necessary to work with partners on multiparty problems toward mutually desirable ends.

Gray (1989) identified several benefits that arise when collaboration of key stakeholders takes place to address problems that cut across public and private sector interests. First, collaboration increases the quality of the solutions, such as management plans considered by the parties, because the collective capacity to respond to the problem is increased. Second, early participation in a multiparty collaboration can minimize the possibility of conflict that might occur during the later stages of a project. Third, the process of collaborating provides certain guarantees that each party’s interests will be protected. Finally, collaboration builds a sense of ownership of the problem that will raise the level of commitment to implementing the solution.

Because ecosystem management is almost invariably a transboundary, multi-party approach to solving “meta-problems,” participation and collaboration of key stakeholders is widely viewed as the single most important element of a successful outcome (Grumbine 1994; Westley 1995; Duane 1997). In a comprehensive survey of ecosystem management initiatives, Yaffee *et al.*

Evaluating Ecosystem Management Capabilities

(1996) found that collaboration of key stakeholders was the single most important factor which enabled projects to reach a quality outcome. Specifically, collaboration within and among public agencies and businesses was an important mechanism for increasing cooperation and communication, fostering trust, and allowing for a more effective outcome that met a greater set of interests. Innes (1996) examined the role of consensus building through case studies of environmental problems involving multiple issues that cut across jurisdictional boundaries. All the cases involved shared power across agencies and levels of government, and between private and public sectors. Innes found collaboration not only increased trust, communication and the development of public-private networks, but also resulted in stronger outcomes or plans that were beneficial to the resource or to the natural system as a whole. More recently, Kennedy *et al.* (2000) found in their analysis of 100 cases involving watershed management in the U.S. that collaboration by stakeholders was a key feature in improving resource management.

In Florida, it is largely recognized at the state level that ecosystem approaches to management are an important aspect of effective environmental management. Ecological systems, particularly regional watersheds, extend across multiple jurisdictions making sustainable management of the entire natural system more complicated (Kirklin 1995). Because ecosystems often do not adhere to what has become a “crazy quilt” of land ownership and governance, environmental management goals are not being reached and natural systems such as the Everglades continue to decline (Light *et al.* 1995; Daniels *et al.* 1996). While this natural system is intricately connected over broad spatial and temporal scales, the land use decision-making framework is limited to local jurisdictions and some limited input from regional planning councils. Uncoordinated local land use decisions have cumulative negative impacts on the

Evaluating Ecosystem Management Capabilities

system as a whole. In other words, the Everglades and its sub-ecological systems are suffering a “death” from thousands of locally imperceptible, individual development decisions.

Collaboration across jurisdictional lines and among multiple organizations thus becomes imperative if approaches to ecosystem management are to be attained (Daniels and Walker 1996; Randolph and Bauer 1999). Understanding the statistical and spatial mosaic of management at the ecosystem level is the first step in protecting critical habitats within local planning processes and facilitating collaborative ecosystem management in the future.

Florida is an ideal state to study ecosystem management capabilities at the local level because it requires that each local community prepare a legally binding comprehensive plan. City and county comprehensive plans in Florida stem from the 1985 *Local Government Comprehensive Planning and Land Development Act*, which mandated new local comprehensive plans be written for each local jurisdiction and required that they be consistent with goals of the State plan.

While there are many different types of resource management plans in Florida (e.g., Water Management District plans, Ecosystem Management Area plans, National Estuary Program plan, etc.), comprehensive plans follow a consistent format (in terms of production, element types, and review/updating processes), are an institutionalized policy instrument, and most importantly provide a basis for city and county land use and resource management decisions. Rule 9J-5, adopted by the Department of Community Affairs (DCA) in 1986, requires that specific elements be included in local plans and prescribes methods local governments must use in preparing and submitting plans. Required elements, among others, include land use, coastal management (where applicable), conservation, and intergovernmental coordination. In each element, the rule

Evaluating Ecosystem Management Capabilities

lists the types of data, issues, goals, and objectives that must be addressed using a “checklist” format (May *et al.* 1996). For example, in the conservation element, objectives must conserve wildlife habitat and policies must pursue cooperation with adjacent local governments to protect vegetative communities (9J-5.013). Because these plans must look beyond jurisdictional boundaries, drive collaborative efforts with other jurisdictions or organizations, and contain policies that seek to protect critical habitats comprising broader ecosystems, they act as strong gauges of how well local jurisdictions will manage ecosystems over the long term.

Conceptualizing Ecosystem Plan Quality

To evaluate the ability of local plans to integrate the principles of collaborative ecosystem management, we constructed a coding protocol based on two major components of plan quality. A total of 58 indicators within the components serve to operationalize and measure the degree to which local comprehensive plans in Florida are managing natural systems that traverse multiple jurisdictions. A complete listing of ecosystem management indicators by component and category are provided in Table 1.

The first component is termed *Inter-organizational Coordination and Capabilities*, which captures the ability of a local jurisdiction to collaborate with neighboring jurisdictions and organizations to manage what are often transboundary natural resources. This is a key component in defining local ecosystem plan quality because it measures the degree to which a local community is able to recognize the transboundary nature of natural systems in Florida and coordinate with other parties both within and outside of its jurisdictional lines. An

Evaluating Ecosystem Management Capabilities

intergovernmental coordination plan element is required by the state of Florida, but there is wide variation among plans with regard to protecting natural systems. This plan quality component addresses the critical factors necessary to foster collaboration which include, among other indicators, coordination with public and private entities, joint fact finding, information sharing, inter-governmental agreements, conflict management processes, and integration of other regional environmental plans in the region. In addition, joint database production, cooperative agreements, and the commitment of financial resources indicate that power and information are being shared among individuals and organizations. These indicators are also a sign of trust and reciprocity formed through informal relationships and interpersonal networks.

The second component, *Policies, Tools & Strategies*, represents the heart of a plan because it actualizes community goals and objectives by setting forth actions to protect critical habitats and related natural systems. This plan component draws heavily on the environmental and land-use planning literatures to identify tools that effectively protect ecological systems and their components (Duerksen *et al.* 1997). Policies include traditional regulatory tools, such as land use or density restrictions, control or removal of exotic species, buffer requirements, and the maintenance of wildlife corridors. Incentive-based tools that encourage landowners to protect critical ecological components as opposed to making them do so are another category of indicators in the plan component. They include clustering, density bonuses, the transfer of development rights (TDRs), preferential tax treatments, and mitigation banking.

Land acquisition programs are another important category because they indicate the ability of jurisdictions to fund the purchase of critical habitats and sensitive lands. Florida is considered a

Evaluating Ecosystem Management Capabilities

leader in acquisitions efforts across the country (Beatley 2000). Under its Preservation 2000 Initiative, the State generated \$300 million per year for 10 years to fund the acquisition of sensitive lands. However, leadership at the state level has not necessarily translated into local initiatives to acquire areas containing critical habitat. Finally, other indicators are included in this plan component, such as the designation of special taxing districts, control of public investment, and educational efforts on the importance of protecting significant habitats. In particular, educational programs are essential for engaging stakeholders in the planning process and in helping generate a plan that is enduring and enforceable in its implementation.

(Insert Table 1 about here)

We selected four indicators of particular importance from the coding protocol for more detailed analysis. Mapping their spatial distribution across ecosystems highlighted spatial gaps in the fabric of management that could be corrected by future plan updates. The following four indicators provided an opportunity to test the level of spatial coverage for specific strategies and policies.

Actions to Protect Natural Resources Crossing Into Other Jurisdictions: In Florida, environmental land use decisions involving the protection of critical habitats and ecosystem components are made through local level comprehensive plans. To effectively manage ecosystems that extend beyond the boundaries of a single jurisdiction, it is important that multiple cities and counties coordinate their environmental policies. Local jurisdictions must not only recognize the fact that certain ecological communities and watershed systems are

Evaluating Ecosystem Management Capabilities

transboundary, but also take collaborative measures to ensure that critical natural resources are protected by neighboring communities.

Sharing Information With Other Organizations: Information is power; the way it is collected, stored, and disseminated is a crucial part of designing effective approaches to ecosystem management (Grumbine 1994). One of the largest barriers to ecosystem management is not acquiring enough information, but sharing it across jurisdictional boundaries, agencies, and other organizations (Lee 1992). To this end, data must be widely accessible and highly integrated into all stages of the decision-making process. Joint fact-finding, information networks, data negotiation, and communication can help make certain that information (from both the social and natural sciences) critical to understanding ecosystem-based issues reaches all of the parties involved (Yaffee and Wondolleck 1997). In a recent study investigating collaborative natural resource management involving 30 different initiatives, Schuett *et al.* (2001) found that information exchange among stakeholders was one of the keys to successful collaboration and more effective resource management.

Establishment of Wildlife Corridors: Wildlife corridors are essential landscape components for maintaining the function of natural systems. They provide landscape connectivity for wildlife movement and provide stepping-stones to keep species matrices functioning and intact (Noss 1983; Van Lier and Cook 1994). In cases where a reserve or “patch” is not large enough to sustain a wide-ranging species, corridors may provide a means for increasing the range by creating a network of connected patches. Equally important, they provide a route for wildlife movement, population dispersal and genetic variability among species (Noss and Cooperider

Evaluating Ecosystem Management Capabilities

1994). River systems are examples of natural corridors that maintain aquatic conditions such as water temperature and oxygen content.

One of the major problems facing natural resource planners is that urban development blocks or disrupts natural corridors. Fragmenting corridors can: 1) reduce area of habitat available to species; 2) increase likelihood of population extinction by limiting immigration; and 3) exacerbate genetic problems resulting from inbreeding (Dramstead *et al.* 1996; Peck 1998). For these reasons, high quality environmental plans seek to protect and/or create corridors to facilitate connectivity.

Wildlife corridors are particularly important planning tools for protecting ecological systems in Florida due to the presence of mobile species with large home ranges. The Florida Panther (*Puma concolor coryi*), a high-visibility species historically found in south Florida, has been recorded as having a dispersal range of 233 km from natal range (Maehr *et. al.* 2000).

Other highly mobile species included in the study area considered essential for the protection biodiversity include the Black Bear (*Ursus americanus*), the bobcat (*Lynx rufus*) and other mammalian species. The Florida Fish and Wildlife Conservation Commission identified strategic habitat conservation areas (SHCAs) for all three species in the southwest portion of the State (Cox *et al.* 1994). These habitats occur outside of existing protected areas and may extend across multiple jurisdictions. Corridors that span several jurisdictions may therefore be necessary to facilitate the movements of key species and maintain their habitat mosaics.

Evaluating Ecosystem Management Capabilities

Controls on Exotic and Invasive Species: The ability of exotics to adapt themselves to a variety of natural conditions, persist in new environments, and disperse long distances allows them to often out compete and displace native species. The establishment of nonnative species can adversely affect ecosystem properties, including productivity, water flows, nutrient cycling, and disturbance regimes (Vitousek 1986). Exotics negatively impacting local ecosystems require special attention in planning and management (Peck 1998). Invasive species pose particular problems in Florida where the USGS estimates persistence of 49 aquatic plant and 273 wildlife species alone, ranging from Asian Marshweed (*Limnophila sessiliflora*) and Eurasian Water-milfoil (*Myriophyllum spicatum*) to the Asian Clam (*Corbicula fluminea*) and the Greenhouse Frog (*Eleutherodactylus planirostris*) (USGS 2003; FFWCC 2003). Planning efforts to control these and other potentially destructive species include eradication programs, planting restrictions, and monitoring of disturbed areas.

Methods and Data Analysis

Sample Selection

Sixteen adjacent ecosystem management areas (EMAs) defined primarily by watershed boundaries were selected for analysis in the southern portion of Florida (Figure 1). These ecosystems stretch from the west coast near Tampa Bay to the heavily developed southeast coast of the State, representing a wide variety of biophysical regions and institutional/political settings. A sample of local jurisdictions was then selected from among those jurisdictions containing land area within one of the sixteen EMAs. All counties intersecting the EMAs, plus the fifteen largest

Evaluating Ecosystem Management Capabilities

cities in land area (since the goal is to achieve the greatest level of spatial coverage, cities were selected based on area rather than by population) were selected to form a sample of 45 adjacent local jurisdictions (Figure 2). On average, there are 6.35 jurisdictions within an EMA.

The most recent comprehensive plans for these counties and cities were evaluated against the ecosystem planning protocol containing the indicators to determine their collective ability to manage EMAs. Plans were evaluated by two trained coders working independently of each other. An “inter-coder reliability score” was computed equal to the number of coder agreements for indicators divided by the total number of indicators. We calculated a score of 97%. The literature suggests that an inter-coder reliability score in the range of 90% is generally considered acceptable (Miles and Huberman 1984).

(Insert figure 1 and 2 about here)

Concept Measurement

The Florida Department of Environmental Protection (DEP) defines EMAs and their boundaries based primarily on hydrological connections (water flow being the major landscape component defining ecosystems in Florida), but also take the following additional criteria into account: 1) *Conservation Lands* – existing conservation lands and vegetation community types that do not adhere to watershed boundaries were incorporated into ecosystem management units; 2) *Human Uses and Impacts* – because human uses and associated impacts in an area dictate the type of

Evaluating Ecosystem Management Capabilities

management activities needed, areas with similar uses were integrated into EMAs; 3) *Agency Jurisdiction* – to minimize the degree of overlap within major governmental jurisdictions, DEP District Offices and Water Management Districts were incorporated into EMAs; and 4) *Overall Manageability* – to increase the effectiveness of managing ecological systems, the overall size and complexity of EMAs designations were considered (DEP 1995).

The comprehensive plan for each jurisdiction located within a selected EMA was evaluated against the 58 ecosystem management indicators (Table 1). If a policy or coordination strategy in a plan was mandatory (using the words “must,” “shall,” or “will”), it was coded as a 1. If the indicator was either suggested, faintly present, or not present at all, it was coded as a 0. With this method, the spatial coverage for each indicator could be mapped and measured across the sample of EMAs, spatially identifying gaps in the regional framework of management across southern Florida.

We measured the percentage of spatial coverage for each indicator within each EMA in two stages. First, we computed the proportion (P_{ij}) of the areas in the i^{th} EMA that was occupied by the j^{th} jurisdiction. Second, we used this proportion to weight that jurisdiction’s contribution to the EMA’s score on that indicator (I_{ijk}). That is,

$$P_{ik} = \sum_j P_{ij} I_{ijk}.$$

Next, we computed a total ecosystem plan quality score ($\sum EPQ$) score for each EMA in a similar manner. Specifically, we summed across all of the indicators for a given jurisdiction’s plan and

Evaluating Ecosystem Management Capabilities

normalized this score by dividing by the total number of indicators and multiplying by 10 to place the variable on a 0-10 scale (as previously done by Brody, forthcoming). Then, we weighted each jurisdiction's ΣEPQ score by the proportion (P_{ij}) of the EMA's area occupied by that jurisdiction and summed across all jurisdictions within that EMA. That is,

$$\sum EPQ_i = \sum_j P_{ij} \frac{10}{58} \sum_k I_{ijk}$$

Data Analysis

Once plan quality scores for each EMA was calculated and entered into a GIS database, we conducted the following analysis. First, we used a visual and statistical gap analysis of ΣEPQ scores to reveal the mosaic of management across ecosystems in Florida. Specifically, we mapped each of the 58 indicators and total ecosystem management scores (ΣEPQ) to facilitate an examination of both gaps in protection and spatial consistency of policies at a broad scale. Second, we investigated the spatial association of total ecosystem planning scores across the study area to determine if there is a clustered pattern of strong or weak ecosystem management capabilities. We used a Joint Count Statistic and Global Moran's I to form an overall picture of the degree of spatial dependency across the study area. We then employed a local indicator of spatial autocorrelation (LISA) (Anselin 1995) to identify and map specific areas of spatial clustering. LISAs detect significant spatial clustering around individual locations and pinpoint areas that contribute most to an overall pattern of spatial dependence. This technique offered a finer focus to uncover important features or characteristics in explaining ecosystem management capabilities at the local level.

Results

Overall Pattern of Spatial Coverage

Tables 2 and 3 report the percentage of spatial coverage of detailed or mandatory ecosystem indicators for each EMA and across the entire study area. An indicator with a total spatial coverage of below 20% represents an insufficient degree of consistency between multiple jurisdictions and a potential gap in the ecosystem management framework across southern Florida. With respect to the *Inter-organizational Coordination and Capabilities Plan Component* (Table 2), financial commitment to ensure implementation of collaborative strategies is covered by just over 1% of the study area. Collaboration between science and policy organizations (an essential aspect of managing ecological systems) receives less than 6% of total coverage, concentrated primarily in south Florida EMAs. In addition, intergovernmental agreements to protect the integrity of ecosystems cover approximately 13% of the study area. Some of the highest levels of coverage for this indicator occur in South Florida/Kissimmee River (50.72%) and Greater Tampa Bay (36.73%) EMAs where formal regional agreements, such as a National Estuary Program (NEP) are already in place. Finally, participation in ecosystem-based initiatives is specified in the plans of approximately 16% of the sample of local jurisdictions. Many large ecosystem programs, such as NEPs and EMA planning initiatives rely on the participation of cities and counties for successful implementation. The greatest intent to participate from a spatial perspective comes from the South Florida EMAs most directly associated with the Everglades region. In contrast to the weaknesses in the collaborative management framework across the study area, nearly 92% of the sample specifies coordination

Evaluating Ecosystem Management Capabilities

with Water Management Districts and approximately 80% of all jurisdictions integrate other regional plans and policies into their local comprehensive plans.

(Insert Table 2 about here)

Within the *Policies, Tools & Strategies* plan component (Table 3), two important policies for protecting ecosystem components that are absent from all plans in the sample are phasing of development to reduce wildlife disturbance and the designation of urban growth boundaries that do not include critical habitat (Deurksen *et al.* 1994). Policies associated with public funding strategies, such as controlling investment for public projects and capital improvements programming to protect ecosystem components also receive weak mandatory coverage across the study area. Interestingly, EMAs to the north and west of Lake Okeechobee contain the highest concentrations of public funding policies. Finally, mandatory tax-based policies that encourage development patterns that protect critical habitats and ecosystem processes cover less than 20% of the study area. Policies involving preferential tax treatments to protect critical habitats and the designation of special taxing districts to raise funds for land acquisition are found almost entirely in the South Florida Loxahatchee/Hungry Slough and Everglades Agricultural EMAs. It is not surprising that some of the most intense development and private investment in the State occur in these areas.

The highest level of spatial coverage and consistency across EMAs is associated with more traditional land use policies, such as use restrictions in and around critical habitats, protection of endangered and threatened species, and the protection of native vegetation. All of these types of

Evaluating Ecosystem Management Capabilities

policies are readily incorporated into local plans in part because they receive less opposition than other more complicated or untested planning tools. They are also easy to implement and enforce.

(Insert Table 3 about here)

Spatial Analysis of Specific Ecosystem Management Indicators

While examining the spatial coverage of ecosystem management indicators across the entire study area provides a general idea of policy gaps or weaknesses, it does not indicate the degree of coordination within the EMAs themselves. We therefore selected four indicators for a more detailed examination of their spatial distribution. Specific coordination capabilities and policies in the sample of comprehensive plans were mapped and analyzed according to their respective EMAs. Through these four examples, we could more precisely identify deficiencies in the spatial coverage of important ecosystem management strategies across multiple local jurisdictions. The examples also demonstrate the effectiveness of using GIS techniques to assess ecosystem management capabilities at the local level and identify area specific gaps in management at the ecosystem level.

Actions to Protect Natural Resources Crossing Into Other Jurisdictions

As discussed above, collaboration across political and jurisdictional lines is an essential aspect of managing large ecological systems such as watersheds (Wondolleck and Yaffee 2000, Cortner and Moote 1999; Yaffee *et al.* 1996; Randolph and Bauer 1999). Mapping jurisdictions that

Evaluating Ecosystem Management Capabilities

have policies to coordinate with their neighbors and neighbor's neighbors within each EMA reveals gaps in the management framework at the watershed level. For example, in the Great Tampa Bay Area EMA, Pinellas, St. Petersburg, Pasco, and Manatee all have mandatory collaborative policies to help protect ecosystem components in neighboring jurisdictions, while Hillsborough and Polk counties do not have such a policy in their comprehensive plan (Figure 3). The Tampa Bay Watershed is a well-defined ecological system that has a National Estuary Program plan in place. However, it is equally important that local jurisdictions not only participate in the ecosystem plan, but also incorporate specific policies into their legally binding land use instruments that require government to realize that natural systems do not adhere to a single political or administrative entity. Similarly, within the Southwest Coast EMA, Lee and Monroe Counties have policies to protect resources crossing into adjacent jurisdictions, but such policies are absent in the plans of Hendry, Collier, and Broward Counties. In both instances, mapping this indicator by EMA shows where local and regional planners need to set local policies to attain a more complete spatial coverage of riverine and coastal ecological communities.

(Insert Figure 3 about here)

Sharing Information With Other Organizations

Sharing information is an important aspect of coordinating environmental policies among multiple governmental units. Table 2 shows that information sharing covers almost 50% of the study area, but mapping this indicator shows important spatial gaps within several EMAs (Figure 4). For example, jurisdictions in the northern portion of the St. Johns River EMA, (a

Evaluating Ecosystem Management Capabilities

watercourse flowing into Lake Okeechobee), contain detailed commitments to share information pertaining to the management of ecological systems. However, the ecosystem management indicator is missing from the plans of jurisdictions to the south. Furthermore, while the comprehensive plans for Dade and Palm Beach Counties (the anchor jurisdictions for South Florida Lower East Coast and Central Everglades EMAs) contain information sharing strategies, Broward County's plan does not. The absence of detailed information sharing policies for Broward thus acts as significant policy gap and potential barrier for the effective flow of information from one jurisdiction or government organization to the next. If Broward County adopts information sharing policies in the next update of their plan, the exchange of vital ecological data, collaboration between jurisdictions, and the management of the ecosystems could be significantly enhanced.

(Insert Figure 4 about here)

Establishment of Wildlife Corridors

One of the most important concepts originating from the landscape ecology and conservation biology literatures is that habitats do not stand alone, but are connected by the movement of species, water, and natural materials. A spatial analysis of environmental policies illuminates gaps in county and municipal agencies that support the maintenance of corridors. Mandatory policies for the establishment or protection of wildlife corridors are found in the plans of approximately 50% of the study area. Notable policy gaps exist in the central portions of the Upper St. Johns River, South Florida Lower East Coast and Central Everglades EMAs (Figure 5). One of the largest gaps and lack of spatial coverage for corridors, however, occurs in the

Evaluating Ecosystem Management Capabilities

Southwest Coastal EMA. Lee County to the northwest portion of this ecosystem appears to be the only jurisdiction with significant area within the EMA containing mandatory wildlife corridor policies in its plan. This policy gap in and around Collier County potentially leaves panther, black bear, and bobcat unprotected by local comprehensive planning frameworks.

On the other hand, efforts have been made by state and federal organizations to provide natural corridors in Florida. The state Department of Transportation and U.S. Fish and Wildlife Service have constructed faunal underpasses along Interstate Highway 75 linking Fort Lauderdale and Naples to allow panther and other mobile species to move unencumbered along natural corridors (Smith 1993; Foster and Humphrey 1995). While there have been some exceptional initiatives to maintain wildlife corridors, our analysis shows that in southwest Florida, wildlife corridor policies and related projects have not been adequately incorporated into local land use decisions where they may have the most significant impact.

(Insert Figure 5 about here)

Controls on Exotic and Invasive Species

As noted earlier, exotic or nonnative species can greatly alter the composition, structure, and function of ecological systems if left unchecked. Due to the seriousness and general recognition of this environmental problem, local policy controls on exotic and invasive species are covered by almost 70% of the study area (see Table 3). Jurisdictions in the South Florida EMAs south of Lake Okeechobee, where the introduction of exotic species is greater than any other drainage

Evaluating Ecosystem Management Capabilities

area in the State, appear to pay particular attention to the issue in their comprehensive plans. Despite wide coverage of this ecosystem management indicator, important policy gaps do exist (Figure 6). For example, the absence of exotic controls in the Hendry County comprehensive plan results in significant gaps in the Caloosahatchee to Lee Coast and the Southwest Coast EMAs. The Giant Rams-horn Snail (*Marisa cornuarietis*) is just one well-known exotic species observed in Hendry County. This snail is native to northern South America and was most likely released by aquarium hobbyists. Studies show that this species retards the growth of aquatic plants by feeding on the roots of the plants (Fuller and Benson 1999). The lack of controls in Hendry County could be influential in the Snail's establishment in neighboring Palm Beach and Broward Counties.

Polk, Hardee, and De Soto Counties in the western part of the Greater Charlotte Harbor EMA are also missing this important ecosystem planning policy in their local management frameworks. The Cuban Treefrog (*Osteopilus septentrionalis*) has been established in Polk County for at least ten years. This species preys upon smaller native treefrogs and may reduce their populations via competition and predation (Asthon and Asthon 1988). Brazilian Pepper (*Schinus terebinthifolius*) is another exotic species established in Polk and De Soto counties that was originally imported from Brazil as an ornamental in the 1840s. This plant species forms dense thickets of woody stems that can shade out and displace native vegetation. It also produces certain allelopathic agents, which appear to suppress other plant's growth (Mahendra *et al.* 1995). Brazilian Pepper is now estimated to occupy over 283,400 ha in central and south Florida (Ferriter 1997). In general, it is important for adjacent local jurisdictions to implement consistent exotic control strategies since nonnative species can spread rapidly across large areas.

Evaluating Ecosystem Management Capabilities

(Insert Figure 6 about here)

Evaluation of Total Ecosystem Planning Scores

The Σ EPQ score computed from the area-weighted sum of all indicators allowed us to evaluate the overall management capabilities of multiple jurisdictions for each EMA. This phase of analysis serves as a global assessment of the relative strength of ecosystem management. In general, the highest total ecosystem planning scores occur for coastal EMAs containing areas of valued critical natural resources that are high priorities for protection (Figure 7). For example, EMAs covering the well-known Everglades region south of Lake Okeechobee all receive above average scores. The Greater Tampa Bay EMA, a similarly well-known ecosystem with multiple environmental initiatives including a NEP, also scored comparatively high.

The highest scores, however, appear in the South Florida Everglades Agricultural and Loxahatchee /Hungryland Slough EMAs along the southeast coast of the State. The ecosystem planning capabilities for these adjacent EMAs are well above average, primarily due to the strength of the Palm Beach County plan, which is one of the highest scoring jurisdictions among the sample of comprehensive plans. In contrast, inland EMAs with lower levels of biodiversity and lesser-known natural value receive lower scores. The Upper St. John River and South Florida Appattah Flats EMAs are two notable exceptions. Further study and analysis is needed to understand specifically why EMAs in certain locations receive higher scores than others.

Evaluating Ecosystem Management Capabilities

(Insert Figure 7 about here)

Evaluation of Spatial Distribution of Local Ecosystem Planning Scores

While the measure and comparison of ΣEPQ scores for each EMA enabled us to make overall assessments of ecosystem management capabilities, it could not provide detail on areas of high quality management within the EMAs themselves. In the final phase of the descriptive analysis, we used measures of spatial autocorrelation to assess the geographic distribution of ΣEPQ scores. This approach allowed us to determine if there is significant spatial association of plan scores for adjacent jurisdictions across the study area and identify clusters of high or low scores within specific EMAs.

We conducted three tests for spatial autocorrelation: a Joint Count Statistic, a Global Moran's I and a Local Moran's I (Table 4). All three tests indicate statistically significant spatial autocorrelation among local jurisdictions ($p < .05$) as well as a tendency for high ecosystem planning scores to cluster geographically within EMAs. Both the Joint Count and Global Moran's I statistics show a significantly nonrandom or clustered pattern of scores across the entire study area. Mapping the most statistically significant Local Moran's I scores ($p < .05$) enabled us to identify the location of clusters or hot spots of adjacent jurisdictions with high ΣEPQ scores.

(Insert Table 4 about here)

Evaluating Ecosystem Management Capabilities

Figure 8 illustrates high scoring county or city jurisdictions that are surrounded by jurisdictions that also have high EPQ scores. These jurisdictions serve as hot spots of significantly greater capabilities to manage ecological systems over the long term. For example, a particularly strong clustered pattern of high ecosystem planning scores occurs in the Greater Tampa Bay EMA. The comprehensive plan for Pinellas County in the western part of this watershed is distinguished for its emphasis on ecosystem management and the protection of critical habitats. This plan received the highest total score in the sample of jurisdictions. Upon further review of the comprehensive plans within the Greater Tampa Bay EMA, we observed that cities and counties adjacent to Pinellas seem to borrow from its plan environmental data, descriptions of programs, and wording of specific policies. The trend suggests that an extremely high scoring comprehensive plan may influence the quality of plans in neighboring jurisdictions through communication, collaboration, and information sharing. This phenomenon is most noticeable for cities located within a specific county. Another potential factor that seems to contribute to a spatial clustering of high scoring plans is that the Tampa Bay watershed has a history of collaborative environmental projects (most notably a NEP and associated plan), where there are previously established lines of communication, information sharing, and joint data production. The presence of a collaborative network may facilitate the dispersion of plan content, resulting in an intense hot spot of high scoring plans.

(Insert Figure 8 about here)

The same situation occurs on the east coast of the state with the case of Martin County. Martin is the third highest scoring plan in the sample and is adjacent to Palm Beach, which received the

second highest score. Together, these jurisdictions appear to be partly responsible for the high Σ EPQ scores for the Everglades Agricultural Area and Loxahatchee /Hungryland Slough EMAs. The strong strategies, programs, and policies in the Martin plan seem to have diffused to neighboring jurisdictions and increased the quality of their plans. The result is a concentrated hot spot of high quality ecosystem-oriented local plans that contribute to the effective management of a large ecological unit. While mapping Local Moran's I values reveals an important trend from a planning perspective, results need to be interpreted with caution. Florida is a peninsular landmass, so there are relatively more edges and fewer neighbors. Therefore, these findings should be considered preliminary until more data can be gathered and examined on spatial association of planning scores.

Conclusions and Policy Implications

Our results indicate that there are important policy gaps in the local mosaic of management across ecological systems in southern Florida. In fact local jurisdictions must play a vital role in protecting the structure and function of transboundary ecosystems, planners and managers need to consider the overall spatial distribution of policies among multiple counties and cities. Thinking about environmental issues related to place and space that extend beyond a single jurisdiction is thus essential, particularly in a state where local comprehensive plans are the primary legal policy instruments guiding the land use and resource management actions at the local level.

Evaluating Ecosystem Management Capabilities

In general, we found that incentive and financial based strategies, such as preferential tax treatments and special funding programs are lacking in coverage compared to more traditional land use regulatory tools (see also Brody, forthcoming). In terms of the spatial distribution of indicators within specific EMAs, notable gaps persist in the Southwest Coast, Southeast Coast, and Central Everglades EMAs, particularly for wildlife corridors and collaboration with neighboring jurisdictions. By viewing the general pattern of policies across ecosystems, we are able to conclude that if Collier and Broward counties install specific environmental policies in their comprehensive plans, there will be a more consistent and complete management framework for these EMAs. By measuring and mapping Σ EPQ scores, results show that planners should focus ecosystem management efforts on less developed, inland-dominated EMAs. This recommendation would entail a more proactive approach to local planning, where ecosystem protection strategies are established before significant development takes place, as opposed to a traditional reactionary policymaking stance.

Finally, the discovery that high quality plans tend to occur in neighboring jurisdictions, and that there are literally hot spots of strong ecosystem management capabilities has significant policy implications. From a strategic perspective, if state and regional planners know that by boosting the quality of one jurisdiction's plan, it is likely that the plans of surrounding jurisdictions will also be enhanced, they can focus limited time, personnel, and financial resources on a single jurisdiction as a way to improve the management of a larger ecologically defined area. Thus, rather than implementing ecosystem programs across broad regions, state managers may consider pinpointing one or a few jurisdictions as a strategy to achieve regional management.

Evaluating Ecosystem Management Capabilities

In general, using GIS techniques to map, measure, and analyze the existing mosaic of management across ecological systems in southern Florida helps form a clearer picture of how local jurisdictions can join together to protect transboundary critical natural resources. Mapping specific strategies and policies provides a rapid assessment of ecosystem protection and provides a strategic tool with which to plan more effectively at the level of natural systems. By locating the spatial gaps (or policy weaknesses) within layers of planning policies and among groups of aggregated policies, the total resource management system can be better understood and strengthened through future plan updates. Rather than casting a broad regulatory net, a spatially focused approach will be more precise, efficient, and cost effective. These techniques can be extremely useful for state and regional planners interested in managing large ecological units such as watersheds assist local planners who understand protecting their own natural resource base requires focusing beyond their single jurisdiction. We have analyzed just a few of the possible indicators that can be gathered from local plans or other policy instruments. Managers in Florida and other states could build a GIS database where hundreds of policies, strategies, and other types of information could be rapidly analyzed.

While this study makes a first step at introducing techniques to explore the local mosaic of management across selected ecosystems in Florida, more research and data are needed to more thoroughly understand how and where local jurisdictions can collectively protect ecological systems. For example, we describe the existing pattern of policies across the study area and pinpoint specific examples where improvement can be made, but future work needs to examine the factors driving the quality of management, particularly at the ecosystem level.

Socioeconomic, demographic, and biophysical variables may all play a role in explaining the

Evaluating Ecosystem Management Capabilities

ability of multiple jurisdictions to manage larger watersheds. In addition, future studies will need larger samples and more sophisticated techniques for analyzing the presence spatial of autocorrelation to make conclusions about the degree and cause of clustering of plan scores.

Literature Cited

- Anselin, L. 1995. Local indicators of spatial association - LISA. *Geographical Analysis* 27(2), 93-115.
- Asthor, R.E. and P.S. Asthor. 1988. *Handbook of Reptiles and Amphibians of Florida. Part Three: The Amphibians*. Miami, FL: Windward.
- Beatley, T. 2000. Preserving Biodiversity: challenges for planners. *APA Journal*, 66(1), 5-20.
- Blumenthal, D. and Jannink, J.L. (2000). A Classification of Collaborative Management Methods. *Conservation Ecology* 4(2):
- Brody, Samuel D. Implementing the Principles of Ecosystem Management Through Land Use Planning. *Population and Environment*. Forthcoming.
- Brody, Samuel D. Measuring the Effects of Stakeholder Participation on the Quality of Local Plans Based on the Principles of Collaborative Ecosystem Management. *Journal of Planning Education and Research*. Forthcoming.
- Brown, L.D. 1991. Bridging Organizations and Sustainable Development. *Human Relations* 44(8): 807-831.
- Bryson, J.M. and B. Crosby. 1992. *Leadership for the Common Good: Tackling Public Problems in a Shared-Power World*. San Francisco: Jossey-Bass.
- Clark, W.C. 1989. Managing Planet Earth. *Scientific American* 261(3): 46-57.
- Collinge, S. K. 1996. Ecological consequences of habitat fragmentation: Implications for landscape architecture and planning. *Landscape and Urban Planning*, 36(1): 59-77.
- Cooperider, D. and W. Passmore. 1991. Global Social Change: A New Agenda for the Social Sciences. *Human Relations* 44(10): 1037-1055.

Evaluating Ecosystem Management Capabilities

- Cortner, H. and Moote, M. 1999. *The Politics of Ecosystem Management*. Washington, DC: Island Press.
- Cox, James, Kautz, R., MacLaughlin, M., and Gilbert, T. 1994. *Closing the Gaps In Florida's Wildlife Habitat Conservation System*. Tallahassee, FL: Florida Game and Fresh Water Fish Commission.
- Christensen, Norman, A. Bartuska, J. Brown, S. Carpenter, C. D'Antonio, R. Francis, J. Franklin, J. MacMahon, R. Noss, D. Parsons, C. Petterson, M. Turner, and R. Woodmansee. 1996. The Report of the Ecological Society of America Committee on the Scientific Basis for Ecosystem Management, *Ecological Applications* 6(3), pp. 665-691.
- Daniels, S., Lawrence, R., and Alig, R. 1996. Decision-Making and Ecosystem-Based Management: Applying the Vroom-Yetton Model to Public Participation Strategy. *Environmental Impact Assessment Review*, 16, 13-30.
- Daniels, Steven and Walker, G. 1996. Collaborative Learning: Improving Public Deliberation in Ecosystem-Based Management. *Environmental Impact Assessment Review*, 16, 71-102.
- Department of Environmental Protection, Florida. 1995. *Ecosystem Management Implementation Strategy, Action Plan*. Tallahassee, FL: DEP.
- Dramstad, Wenche E., J. Olson, and R. Forman. 1996. *Landscape Ecology Principles in Landscape Architecture and Land-Use Planning*. Washington, D.C.: Island Press.
- Duerksen, C. D. Elliot, N., Thompson, E., and Miller, J. 1997. *Habitat Protection Planning: Where the Wild Things Are*. APA Planning Advisory Report No. 470/471.
- Emery, F. and E. Trist. 1965. The Causal Texture of Organizational Environments. *Human Relations* 18: 21-35.

Evaluating Ecosystem Management Capabilities

- Endter-Wada, Joanna, Blahna, D., Krannich, R., and Brunson, M. 1998. A Framework for Understanding Social Science Contributions to Ecosystem Management. *Ecological Applications*, 8(3), 891-904.
- Ferrister, A., (ed.). 1997. *Brazilian Pepper Management Plan for Florida*. West Palm Beach: Florida EPPC.
- Florida Fish and Wildlife Conservation Commission. 2003. Florida's Exotic Wildlife. <http://wld.fwcc.state.fl.us/critters/exotics/exotics.asp> .
- Foster, M.L. and Humphrey, S. R. 1995. Use of underpasses by Florida Panthers and other wildlife. *Wildlife Society Bulletin*, 23: 95-100.
- Fuller, Pam and Amy Benson. 1999. Nonindigenous Species Introduced into South Florida. Poster presented at the South Florida Restoration Science Forum, May 17-19, Boca Raton, FL.
- Gray, B. 1989. *Collaborating: Finding Common Ground for Multiparty Problems*. San Francisco, CA: Jossey-Bass.
- Gray, B. and D. Wood. 1991. Collaborative Alliances: Moving From Theory to Practice. *Journal of Applied Behavioral Sciences* 27(1): 3-22.
- Grumbine, Edward. 1994. What Is Ecosystem Management? *Conservation Biology*, 8(1), 27-38.
- Innes, Judith. 1996. Planning Through Consensus Building: A New View of the Comprehensive Planning Ideal. *Journal of American Planning Association* 62: 460.
- Kennedy, D.S., S.T. McAllister, W.H. Caile, and J.S. Peckham. 2000. The New Watershed Source Book. Natural Resources Law Center, University of Colorado School of Law: Boulder, CO.

Evaluating Ecosystem Management Capabilities

- Kirklin, John. 1995. Protecting Species and Ecosystems Within Planning Processes. *Environmental Planning*, 12(4), 6-13.
- Lee, R.G. 1992. Ecologically Effective Social Organization As A Requirement for Sustaining Watershed Ecosystems. In R.J. Naiman (ed.), *Watershed Management: Balancing Sustainability and Environmental Change*. New York: Springer-Verlag.
- Light, S., L. Gunderson, and C.S. Holling. 1995. The Everglades: Evolution of Management in a Turbulent Ecosystem. In Lance Gunderson, C.S. Holling, and Stephen S. Light (eds.). *Barriers and Bridges to the Renewal of Ecosystems and Institutions*, pp. 3-34. New York: Columbia University Press.
- Mahendra, K.J., Yu, B., Rogers, J.M., Smith, A.E., Boger, E.T.A., Ostrander, R.L. 1995. Specific competitive inhibitor of secreted phospholase A₂ from berries of *Schinus terebinthifolius*. *Phytochemistry* 39:537-47.
- May, Peter, Raymond Burby, Neil Erickson, John Handmer, Jennifer Dixon, Sarah Michaels, and D. Smith. 1996. *Environmental Management and Governance: Intergovernmental Approaches to Hazards and Governance* (London, Routledge).
- McGinnis, M., Woolley, J., and Gamman, J. 1999. Bioregional Conflict Resolution: Rebuilding Community in Watershed Planning and Organizing. *Environmental Management*, 24(1), 1-12.
- Mehr, David S., E. Darrell Landb, David B. Shindleb, Oron L. Bassc and Thomas S. Hochtord. 2000. Florida panther dispersal and conservation. *Biological Conservation*, 106(2): 187-197.
- Miles, M. and A.M. Huberman. 1984. *Qualitative Data Analysis: A Sourcebook of New Methods*. Newbury Park, CA: Sage.

Evaluating Ecosystem Management Capabilities


- Noss, Reed F. 1983. A Regional Landscape Approach to Maintain Diversity. *BioScience* 33: 700-706.
- Noss, Reed and Allen Cooperrider. 1994. *Saving Nature's Legacy: Protecting and Restoring Biodiversity*, (Washington, D.C., Island Press).
- Noss, Reed and Scott, M. 1997. Ecosystem Protection and Restoration: The Core of Ecosystem Management. In M. Boyce and A. Hanley, (Eds.), *Ecosystem Management: Applications for Sustainable Forest and Wildlife Resources* (pp. 239-264). New Haven: Yale University Press.
- Peck, Sheila. 1998. *Planning for Biodiversity: Issues and Examples*. Washington, D.C.: Island Press.
- Randolph, John and Bauer, M. 1999. Improving Environmental Decision-Making Through Collaborative Methods. *Policy Studies Review*, 16: 169-191.
- Scheutt, Michael, Selin, Steve, and Carr, Deborah. 2001. Making It Work: Keys to Successful Collaboration in Natural Resource Management. *Environmental Management* 27(4): 587-593.
- Selin, Steve and Carr, Deborah. (2000). Modeling Stakeholder Perception of Collaborative Initiative Effectiveness. *Society and Natural Resources* 13: 735-745.
- Smith, D.S. 1993. Greenway Case Studies. In Smith, D.S. and P.C. Hellmund, eds., *Ecology of Greenways. Design and Function of Linear Conservation Areas*. U. of Michigan Press, Minneapolis, pp. 161-208.
- Szaro, R., Sexton, W., and Malone, C. 1998. The emergence of ecosystem management as a tool for meeting people's needs and sustaining ecosystems. *Landscape and Urban Planning*, 40, 1-7.
- USGS. (2003). Nonindigenous Aquatic Plant Records.
<http://nas.er.usgs.gov/queries/plants/PlantStateResults.asp> .

Evaluating Ecosystem Management Capabilities

- Vitousek, P.M. 1986. Biological Invasions and Ecosystem Properties: Can Species Make a Difference? In Mooney and Drake, eds., *Ecology of Biological Invasions of North America and Hawaii*. Springer-Verlag, New York. 163-176.
- Van Lier, Hubert and Edward Cook. (1994). Ecological Networks: a conspectus. In *Landscape Planning and Ecological Networks*. Amsterdam: Elsevier Science.
- Vogt, John, C. Gordon, J. Wargo, and D. Vogt. (1997) *Ecosystems* (NY, Springer-Verlag).
- Westley, Frances. 1995. Governing Design: The Management of Social Systems and Ecosystems Management. In *Barriers and Bridges to the Renewal of Ecosystems and Institutions*, pp. 391-427. Edited by Lance Gunderson, C.S. Holling, and Stephen S. Light. Columbia University Press, New York.
- Westley, Frances and Harrie Vrendenburg. (1997). Interorganizational Collaboration and the Preservation of Global Biodiversity. *Organizational Science* 8(4): 381- 403.
- Wondolleck, Julia and Yaffee, S. (2000). *Making Collaboration Work: Lessons from Innovation in Natural Resource Management*. Washington, D.C.: Island Press.
- Yaffee, S., A. Phillips, I. Frentz, P. Hardy, S. Maleki, and Thorpe, B. (1996). *Ecosystem Management in the United States: An Assessment of Current Experience*. Washington, D.C.: Island Press.
- Yaffee, Steven and Julia Wondolleck. (1997). Building Bridges Across Agency Boundaries. In *Creating A Forestry for the 21st Century*. Edited by Kathryn A. Kohm and Jerry F. Franklin. Washington, D.C.: Island Press.

Figure 1: Ecosystem Management Areas


Evaluating Ecosystem Management Capabilities



Evaluating Ecosystem Management Capabilities

Figure 2: Local Jurisdictions Within the Boundaries of EMAs

Evaluating Ecosystem Management Capabilities



Evaluating Ecosystem Management Capabilities

Table 1: Plan Coding Protocol

Inter-organizational Coordination & Capabilities for Ecosystem Management		
Other organizations/stakeholders identified	Coordination with other organizations/jurisdictions specified	Coordination within jurisdiction specified
Coordination with adjacent counties specified	Coordination with state level entities specified	Coordination with Federal level entities specified
Intergovernmental bodies specified	Joint database production	Coordination with private sector
Information sharing	Links between science and policy specified	Position of jurisdiction within bioregion specified
Intergovernmental agreements	Conflict management processes	Commitment of financial resources
Coordination with Water Management Districts	Participation in Ecosystem-based initiatives (i.e. NEP, EMAs)	Integration with other plans & policies
Other forms of coordination		
Policies, Tools, & Strategies		
<i>A. Regulatory Tools</i>		
Resource use restrictions	Density restrictions	Restrictions on native vegetation removal
Removal of exotic/invasive species	Buffer requirements	Fencing controls
Public or vehicular access restrictions	Phasing of development	Controls on construction
Conservation zones/overlay districts	Performance zoning	Subdivision standards
Protected areas/sanctuaries	Urban growth boundaries to exclude habitat	Targeted growth away from habitat
Establishment of a network of protected areas	Create/maintain wildlife corridors	Protect threatened/endangered species
Capital improvements programming	Site plan review	Habitat restoration actions
Actions to protect resources in other jurisdictions	Structural or design solutions to protect habitat	
<i>B. Incentive-based Tools</i>		
Density bonuses	Clustering away from habitats	Transfer of development rights
Preferential tax treatments	Mitigation banking	Other incentive-based tools
Impact fees to protect habitat	Specific mitigation measures to protect habitat	Other regulatory tools
<i>C. Land Acquisition Programs</i>		
Fee simple purchase	Conservation easements	Other land acquisition techniques
<i>D. Other Policies, Tools, Strategies</i>		
Control of public investments and Projects	Public education programs	Designation of special Taxing districts to raise funds for land acquisition
Studies or ecological surveys		

Evaluating Ecosystem Management Capabilities

Table 2: Ecosystem Management Indicators for Inter-organizational Coordination & Capabilities

POLICIES	UPPER ST. JOHNS RIVER	INDIAN RIVER LAGOON	S. FLORIDA KISSIMMEE RIVER	LAKE WALES RIDGE	GREATER TAMPA BAY	GREATER CHARLOTTE HARBOR	S. FLORIDA ALLAPATTAH FLATS	SARASOTA BAY	S. FLORIDA FISHEATING CREEK	S. FLORIDA TAYLOR CREEK	S. FLORIDA LOXAHATCHEE/HUNGRYLAND SLOUGH	S. FLORIDA EVERGLADES AGRICULTURAL	CALOOSAHATCHEE TO LEE COAST	SOUTHWEST COAST	S. FLORIDA LOWER EAST COAST	S. FLORIDA CENTRAL EVERGLADES	AVG Policy coverage
Commitment of financial resources	9.27	0.00	0.91	5.35	8.48	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.96	0.00	1.62
Link betw. Science and policy	0.00	0.34	0.00	0.00	8.48	0.00	30.92	0.00	0.00	14.31	32.75	0.38	0.00	0.00	0.00	0.00	5.45
Position in ecosystem ID'd	13.97	0.00	14.27	0.00	14.06	14.54	0.00	0.00	1.01	0.00	0.00	0.00	18.76	1.41	36.74	45.14	9.99
IGAs specified	38.60	18.01	50.72	2.83	36.73	8.01	0.29	25.59	0.00	0.00	0.00	0.00	6.65	13.28	0.00	12.85	13.35
Participation in Ecosyst. initiatives	32.38	42.33	0.00	0.00	8.48	0.00	0.00	0.00	0.00	0.00	64.82	84.54	0.00	0.00	18.79	10.06	16.34
Joint database production	24.68	9.34	18.00	57.52	18.21	36.50	30.92	0.00	1.01	14.31	40.08	0.38	12.47	55.75	2.16	0.00	20.08
Intergovernmental bodies	31.80	19.68	13.20	0.00	36.37	21.86	95.84	15.93	1.01	19.79	97.56	84.92	18.76	0.00	18.79	10.06	30.35
Coord. With private sector	35.25	21.42	55.51	8.17	19.82	46.20	10.23	67.07	1.01	79.71	72.14	84.54	12.11	0.00	18.99	10.06	33.89
Other Coord.	54.67	64.10	0.00	0.00	30.94	26.57	39.07	15.93	0.00	14.31	97.56	84.92	7.01	69.03	23.03	22.90	34.38
Coord. Within jurisdiction	17.73	21.77	18.00	57.52	35.85	48.54	39.07	15.93	0.00	14.31	100.00	85.16	0.00	3.21	83.16	80.59	39.11
Coord. With federal	37.41	28.35	30.36	45.01	64.38	17.40	65.21	9.66	99.76	5.47	7.33	2.69	66.58	25.13	66.64	83.38	40.92
Conflict management processes	25.35	18.35	18.00	57.52	41.79	30.04	31.21	15.93	0.00	14.31	97.56	97.19	60.87	40.29	57.81	68.04	42.14
Information sharing	49.57	12.76	32.27	57.52	41.03	30.04	38.78	15.93	0.00	14.31	100.00	84.92	29.39	79.08	55.73	68.04	44.64
Coord. with other Orgs.	66.40	16.27	63.86	47.83	36.87	9.98	95.84	15.93	54.29	19.79	97.56	97.19	60.87	27.01	61.76	55.19	51.67
Other Organizations ID'd	44.86	28.69	52.58	2.83	30.94	21.96	96.13	25.59	45.46	19.79	100.00	99.64	63.10	87.40	62.24	68.04	53.08
Coord. With state	47.85	28.69	30.36	45.01	65.08	49.76	96.13	76.73	99.76	19.79	100.00	87.37	66.58	23.32	62.24	68.04	60.42
Coord. With adjacent jurisdictions	69.19	42.33	100.00	94.65	53.13	81.85	9.94	92.66	98.76	79.71	64.82	99.26	80.03	96.04	57.81	68.04	74.27
Integration with other plans	67.13	70.68	51.33	97.18	98.37	59.73	75.15	92.66	99.76	85.18	72.14	99.26	63.10	87.40	61.96	68.04	78.07
Coord. With WMD's	89.30	71.02	100.00	100.00	75.56	90.92	100.00	76.73	99.77	99.49	100.00	99.88	92.14	97.85	89.70	93.44	92.54

Evaluating Ecosystem Management Capabilities

Table 3: Ecosystem Management Indicators for Policies, Tools, & Strategies

	UPPER ST. JOHNS RIVER	INDIAN RIVER LAGOON	S. FLORIDA KISSIMMEE RIVER	LAKE WALES RIDGE	GREATER TAMPA BAY	GREATER CHARLOTTE HARBOR	S. FLORIDA ALLAPATTAH FLATS	SARASOTA BAY	S. FLORIDA FISHEATING CREEK	S. FLORIDA TAYLOR CREEK	S. FLORIDA LOXAHATCHEE/HUNGRYLANDSLOUGH	S. FLORIDA EVERGLADES AGRICULTURAL	CALOOSAHATCHEE TO LEE COAST	SOUTHWEST COAST	S. FLORIDA LOWER EAST COAST	S. FLORIDA CENTRAL EVERGLADES	AVG Policy coverage
Urban growth boundaries	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Phasing development	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Control public investments	26.24	33.34	0.91	5.35	14.42	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	5.02
Subdivision standards	0.00	0.34	0.00	0.00	0.00	0.00	30.92	0.00	0.00	14.31	32.75	0.38	0.00	13.28	0.00	12.85	6.55
Other incentives	22.97	18.01	0.91	5.35	64.73	33.84	0.29	0.00	1.03	0.00	0.00	0.00	12.11	0.00	0.00	0.00	9.95
Targeted growth areas	21.13	8.99	0.00	0.00	16.88	26.53	0.00	15.93	0.02	0.00	0.00	12.27	25.19	30.25	0.00	12.85	10.63
Capital improvements	9.75	3.41	0.91	5.35	8.48	14.60	0.00	0.00	1.01	0.00	0.00	0.00	41.14	10.04	36.74	45.14	11.04
Fencing controls	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	64.82	84.54	0.00	13.28	18.79	22.90	12.77
Preferential tax treatments	0.00	0.34	0.00	0.00	0.00	0.00	30.92	0.00	0.00	14.31	97.56	84.92	0.00	0.00	18.79	10.06	16.06
Special taxing districts	0.00	0.34	0.00	0.00	8.48	0.00	30.92	0.00	0.00	14.31	97.56	84.92	0.00	0.00	18.79	10.06	16.59
Mitigation banking	14.99	0.00	0.91	5.35	19.49	14.60	0.00	0.00	1.01	0.00	64.82	84.77	41.14	10.45	46.24	35.46	21.20
Network protected areas	32.66	27.00	18.00	57.52	18.21	21.97	0.29	0.00	0.00	0.00	64.82	84.54	0.00	0.00	18.79	10.06	22.12
Impact fees	15.41	16.26	0.00	0.00	0.37	0.69	87.98	0.00	0.00	19.79	97.56	84.92	0.00	13.28	18.79	22.90	23.62
Structural solutions	31.96	33.34	0.91	5.35	8.48	14.54	0.00	0.00	1.01	0.00	64.82	97.05	37.30	18.77	44.25	35.46	24.58
Density bonuses	43.97	8.99	27.42	45.01	47.79	2.70	0.00	0.00	54.29	0.00	64.82	84.54	0.00	0.00	18.79	10.06	25.52
Performance Zoning	56.37	42.67	38.43	8.17	18.34	37.94	30.92	83.00	0.02	14.31	32.75	12.66	54.23	25.60	0.00	0.00	28.46
Other land acquisition	39.95	27.35	1.07	0.00	53.37	19.42	39.07	0.00	0.00	14.31	97.56	84.92	35.68	21.92	18.79	22.90	29.77
Public access controls	17.45	44.01	0.00	0.00	42.63	32.59	87.98	92.66	1.01	19.79	32.75	0.38	48.15	79.08	38.71	57.98	37.20
Fee simple purchase	42.68	27.34	26.51	39.66	42.63	22.34	31.21	83.00	54.29	14.31	104.89	84.92	0.36	55.75	18.99	10.06	41.18

Evaluating Ecosystem Management Capabilities

Table 3: Ecosystem Management Indicators for Policies, Tools, & Strategies

POLICIES	UPPER ST. JOHNS RIVER	INDIAN RIVER LAGOON	S. FLORIDA KISSIMMEE RIVER	LAKE WALES RIDGE	GREATER TAMPA BAY	GREATER CHARLOTTE HARBOR	S. FLORIDA ALLAPATTAH FLATS	SARASOTA BAY	S. FLORIDA FISHEATING CREEK	S. FLORIDA TAYLOR CREEK	S. FLORIDA Loxahatchee/Hungrylands Slough	S. FLORIDA EVERGLADES AGRICULTURAL	CALOOSAATCHEE TO LEE COAST	SOUTHWEST COAST	S. FLORIDA LOWER EAST COAST	S. FLORIDA CENTRAL EVERGLADES	AVG Policy coverage
Protection in other jurs.	54.40	42.67	29.45	39.66	42.30	10.09	30.92	15.93	98.75	14.31	97.56	87.37	47.83	21.92	20.75	22.90	42.30
Protected areas	15.41	9.34	0.00	0.00	31.63	32.59	38.78	83.00	1.01	14.31	97.56	97.19	66.70	82.76	57.50	55.19	42.69
Site plan review	64.51	46.09	39.22	8.17	43.42	27.38	38.78	15.93	44.47	14.31	100.00	99.64	73.38	81.35	23.23	10.06	45.93
Conservation easements	46.88	33.93	15.29	45.01	82.70	41.71	65.21	83.00	54.31	5.47	72.14	84.54	29.39	64.39	20.98	10.06	47.19
Wildlife corridors	48.96	34.28	30.36	45.01	25.07	17.30	96.13	0.00	99.76	19.79	97.56	87.37	59.93	10.04	57.81	55.19	49.03
habitat restoration	55.35	67.60	14.22	45.01	47.88	24.46	88.27	25.59	55.30	19.79	97.56	97.43	43.95	32.06	46.21	48.30	50.56
Clustering	60.45	34.27	30.36	45.01	73.47	59.99	88.27	83.00	98.76	19.79	40.08	15.11	73.38	94.63	0.20	12.85	51.85
Construction contols	48.16	34.28	38.31	2.82	48.64	23.79	96.13	25.59	45.46	19.79	97.56	87.61	66.58	25.13	80.99	93.44	52.14
Conservation zones	70.20	42.67	68.83	100.00	46.60	65.62	30.92	25.59	55.32	14.31	97.56	85.16	12.47	57.56	46.21	35.46	53.41
TDRs	55.16	37.69	45.58	97.18	98.01	58.94	96.13	83.00	55.30	19.79	100.00	84.92	12.47	69.03	18.99	22.90	60.00
Ecological studies	78.26	71.02	39.22	8.17	19.49	40.38	96.13	0.00	44.47	19.79	100.00	99.88	80.03	97.85	81.19	93.44	60.89
Public education programs	74.12	71.01	27.42	45.01	88.28	37.04	88.27	92.66	55.30	19.79	100.00	84.92	47.79	23.32	55.73	68.04	61.47
Buffer requirements	96.88	67.61	82.01	47.83	43.76	76.08	106.07	83.00	55.32	99.49	97.56	97.19	66.33	38.88	18.79	22.90	68.73
Control exotics	35.86	62.03	15.17	39.66	87.52	37.15	96.13	92.66	99.76	19.79	100.00	87.61	66.94	80.88	87.74	93.44	68.90
Density restrictions	94.56	64.09	82.02	100.00	91.34	86.80	31.21	15.93	55.32	14.31	97.56	97.19	66.70	96.04	57.81	68.04	69.93
other reg. Tools	94.54	67.27	83.88	47.83	43.00	59.89	75.15	92.66	98.76	85.18	64.82	99.50	79.67	40.69	46.21	48.30	70.46
Protect native vegetation	97.36	71.02	70.57	8.17	88.28	37.50	100.00	83.00	44.45	99.49	97.56	87.61	48.19	80.88	89.50	93.44	75.19
Endangered/threatened species	71.87	37.69	66.80	47.83	82.71	42.55	96.13	83.00	98.75	19.79	100.00	99.88	80.03	97.85	85.15	93.44	75.22
Mitigation measures	80.23	71.02	47.45	97.18	75.56	90.92	96.13	67.07	99.77	19.79	97.56	99.88	85.49	97.85	82.99	93.44	81.40
Use restrictions	92.23	71.02	82.02	100.00	98.01	97.46	96.13	83.00	55.32	19.79	97.56	97.43	73.34	97.85	84.95	93.44	83.72

Evaluating Ecosystem Management Capabilities

Table 4: Tests for Spatial Autocorrelation for Ecosystem Planning Scores


Statistical Test	Expected	Observed	Z-Value	p-value
<i>Joint-Count*</i>				
1-1	36.87	71.5	0.5672	0.5706
0-0	90.6462	101.0	0.0686	0.9454
1-0	117.984	73.0	-5.4064	0.0000
<i>Moran's I</i>	-0.0096	0.15523	2.2688	0.0233
<i>LISA</i>	-0.0096	0.47	2.17	0.0300
	-0.0096	1.24	3.85	0.0001
	-0.0096	1.39	3.2	0.0014
	-0.0096	1.28	2.29	0.0220
	-0.0096	1.11	2.29	0.0220
	-0.0096	1.11	2.29	0.0220
	-0.0096	1.01	4.13	0.0000
	-0.0096	1.3	4.69	0.0000
	-0.0096	1.8	3.21	0.0013
	-0.0096	0.78	1.99	0.0466

*Total ecosystem planning scores were converted into a dichotomous variable (above or below average) to obtain a measure of spatial autocorrelation using joint count statistics.

Evaluating Ecosystem Management Capabilities

Figure 3: Distribution of Policies for Actions to Protect Natural Resources Crossing Into Adjacent Jurisdictions


Evaluating Ecosystem Management Capabilities



Evaluating Ecosystem Management Capabilities

Figure 4: Distribution of Strategies for Information Sharing

Evaluating Ecosystem Management Capabilities



Evaluating Ecosystem Management Capabilities

Figure 5: Distribution of Policies for to Maintain or Create Wildlife Corridors

Evaluating Ecosystem Management Capabilities





Figure 6: Distribution of Policies to Control or Remove Exotic Species

Evaluating Ecosystem Management Capabilities



Evaluating Ecosystem Management Capabilities

Figure 7: Total Ecosystem Management Scores




Figure 8: Spatial Clusters of High Scoring Plans

Evaluating Ecosystem Management Capabilities

